83103 Avenue 48, Ste.1B #204 Coachella, CA 92236 USA Phone: +1.6268339877 Email: info@cali-bio.com

Product Datasheet

Product Name B-Cell Leukemia/Lymphoma 2 Human Recombinant (–BH2)

Cata No CB501489

Source Escherichia Coli.

Synonyms Apoptosis regulator Bcl-2, BCL2, B-cell CLL/lymphoma 2, Bcl-2.

Description

BCL2 gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Two transcript variants, produced by alternate splicing, differ in their C-terminal ends.

Bcl-2 Des BH2 domain (187-202 residues) Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 202 amino acids 1-186 and 203-218.

The Bcl-2 is expressed as His-Tag fusion protein and purified by proprietary chromatographic techniques.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Purity

Greater than 95.0% as determined by:

- (a) Analysis by RP-HPLC.
- (b) Analysis by SDS-PAGE.

Formulation

The protein contains 10mM Tris-HCL pH-8, 1mM EDTA and 250mM NaCl.

Reconstitution

Suspend Bcl-2 in 100µl of 0.5M Acetic acid, over night at 4°C.

Dilute 10 fold into selected buffer system.

BCL-2 has tendency to form intramolecular disulfide bond, 5mM DTT is recommended in assay buffer.

When running SDS-PAGE gel, 10mM DTT is

recommended.

Stability

Lyophilized Bcl-2 although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution Bcl-2 should be stored at 4°C between 2-7 days and for future use below

-18°C.

For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).

Please prevent freeze-thaw cycles.

Applications

Input marker or positive control (Western Blotting). Function study.